



# **VAR-Based Granger-Causality Test in the Presence of Instabilities**

**Barbara Rossi  
Yiru Wang**

**January 2019**

*Barcelona GSE Working Paper Series*

*Working Paper n° 1083*

# VAR-based Granger-causality Test in the Presence of Instabilities

Barbara Rossi

ICREA Professor at University of Pompeu Fabra

Barcelona Graduate School of Economics, and

CREI

Barcelona, Spain.

`barbara.rossi@upf.edu`

Yiru Wang

University Pompeu Fabra

Barcelona, Spain.

`yiru.wang@upf.edu`

## **Abstract**

In this article, we review Granger-causality tests robust to the presence of instabilities in a Vector Autoregressive framework. We also introduce the `gcrobustvar` command, which illustrates the procedure in Stata. In the presence of instabilities, the Granger-causality robust test is more powerful than the traditional Granger-causality test.

*Keywords:* `gcrobustvar`, Granger-causality, VAR, instability, structural breaks, local projections.

# 1 Introduction

Vector Autoregressive (VAR) models have played an important role in macroeconomic analysis since Sims (1980). A VAR is a multi-equation, multi-variable linear model where each variable is in turn explained by its own lagged values, as well as current and past values of the remaining variables. Compared with a univariate autoregression, VARs provide both a systematic way to capture the rich dynamics in multiple time series as well as a coherent and credible approach to forecasting.

Granger (1969) causality is a useful tool for characterizing the dependence among time series in reduced-form VARs, and Granger-causality test statistics are widely used to examine whether lagged values of one variable help to predict another variable – see Stock and Watson (2001).

However, VAR analyses in macroeconomic data face important practical challenges: economic time-series data are prone to instabilities (see Stock and Watson (1996, 1999, 2003, 2006), Rossi (2013), Clark and McCracken (2006b)) and VARs estimates may be prone to instabilities as well (see Boivin and Giannoni (2006), Kozicki and Tinsley (2001), and Cogley and Sargent (2001, 2005)).

Thus, given the widespread use of VARs and the evidence of instabilities, it is potentially important to allow for changes over time when doing VAR-based statistical inference. As demonstrated in Rossi (2005), statistical tests that are based on stationarity assumptions are invalid in the presence of instabilities. Since the traditional Granger-causality test assumes stationarity, it is not reliable in the presence of instabilities and may lead to incorrect inference.

In this article, we present the `gcrobustvar` command, which illustrates how to test Granger-causality in a way that is robust to the presence of instabilities. The test is based on methodologies developed by Rossi (2005) and includes the robust versions of the mean and exponential Wald tests (Andrews and Ploberger (1994)), the Nyblom (1989) test, and the Quandt (1960) and Andrews (1993) quasi-likelihood-ratio test, jointly testing for both parameter instability and Granger-causality. In the presence of instabilities, the Granger-causality robust tests are more powerful than the traditional Granger-causality test. The tests can also be used to find the point in time in which Granger-causality either appears or breaks down in the data. Besides, the test is valid for reduced-form VAR models as well as VAR-based direct multistep (VAR-LP) forecasting models. The former assume homoskedastic idiosyncratic shocks, while the latter are estimated via Local Projections (see Jordà (2005)), and, hence, assume heteroskedastic and serially correlated idiosyncratic shocks.

We first introduce the tests, then present the Stata commands that implement them. Then, we illustrate the empirical implementation of the robust Granger-causality tests using a three-variable (inflation, unemployment and interest rate) VAR model with four lags as in Stock and Watson (2001), as well as a direct multistep VAR-LP forecasting model. Finally, we compare the results with those based on a traditional Granger-causality test.

The remainder of this paper is organized as follows. Section 2 describes the theoretical framework and the Granger-causality robust tests. Section 3 introduces the `grobustvar` command, which implements the Granger-causality robust tests in Stata. Section 4 applies the Granger-causality robust tests in the three-variable VAR and compares the results with the traditional Granger-causality test. Section 5 applies the Granger-causality robust tests in the direct multistep VAR-LP forecasting model.

## 2 VAR-based Granger-Causality Test in the Presence of Instabilities

### 2.1 Framework

We consider two types of VAR specifications. The first is a traditional reduced-form VAR:

$$\begin{aligned}
 A(L)y_t &= u_t \\
 A(L) &= I - A_1L - A_2L^2 - \dots - A_pL^p \\
 u_t &\overset{i.i.d.}{\sim} (O, \Sigma)
 \end{aligned} \tag{1}$$

where  $y_t = [y_{1,t}, y_{2,t}, \dots, y_{n,t}]'$  is an  $(n \times 1)$  vector, and  $A_j, j = 1, \dots, p$ , are  $(n \times n)$  coefficient matrices.

The second is a direct multistep VAR-LP forecasting model. By iterating eq (1),  $y_{t+h}$  can be projected onto the linear space generated by  $(y_{t-1}, y_{t-2}, \dots, y_{t-p})'$ , specifically

$$y_{t+h} = \Phi_1 y_{t-1} + \Phi_2 y_{t-2} + \dots + \Phi_p y_{t-p} + \epsilon_{t+h} \tag{2}$$

where  $\Phi_j, j = 1, \dots, p$  are functions of  $A_j, j = 1, \dots, p$  in eq (1), and  $\epsilon_{t+h}$  is a moving average of the errors  $u$  from time  $t$  to  $t+h$  in eq (1) and therefore uncorrelated with the regressors but serially correlated itself<sup>1</sup>.

Note that  $h = 0$  is a special case where eq (2) degenerates to eq (1), thus we focus on eq (2) from now

---

<sup>1</sup>See Jorda (2005) for more details of Local Projections.

onwards.

To consider a more concrete example, Stock and Watson (2001) study a three-variable VAR with four lags and  $h = 0$ . The variables included are inflation ( $\pi_t$ ), unemployment ( $u_t$ ) and interest rate ( $R_t$ ). Their reduced-form VAR is:

$$\begin{bmatrix} \pi_t \\ u_t \\ R_t \end{bmatrix} = \Phi_1 \begin{bmatrix} \pi_{t-1} \\ u_{t-1} \\ R_{t-1} \end{bmatrix} + \Phi_2 \begin{bmatrix} \pi_{t-2} \\ u_{t-2} \\ R_{t-2} \end{bmatrix} + \Phi_3 \begin{bmatrix} \pi_{t-3} \\ u_{t-3} \\ R_{t-3} \end{bmatrix} + \Phi_4 \begin{bmatrix} \pi_{t-4} \\ u_{t-4} \\ R_{t-4} \end{bmatrix} + \begin{bmatrix} \epsilon_t^\pi \\ \epsilon_t^u \\ \epsilon_t^R \end{bmatrix} \quad (3)$$

$$\Phi_j = \begin{bmatrix} \phi_j^{\pi,\pi} & \phi_j^{\pi,u} & \phi_j^{\pi,R} \\ \phi_j^{u,\pi} & \phi_j^{u,u} & \phi_j^{u,R} \\ \phi_j^{R,\pi} & \phi_j^{R,u} & \phi_j^{R,R} \end{bmatrix}, \quad j = 1, \dots, 4$$

Thus, in Stock and Watson (2001), the reduced-form VAR involves three equations: current unemployment as a function of past values of unemployment, inflation and the interest rate; current inflation as a function of past values of inflation, unemployment and the interest rate; and current interest rate as a function of past values of inflation, unemployment and the interest rate. Stock and Watson (2001) consider traditional Granger-causality tests in each equation where the null hypothesis is:  $H_0^* : \theta = 0$ , where  $\theta$  is the appropriate subset of  $vec(\Phi_1, \Phi_2, \dots, \Phi_p)$ . For example, unemployment doesn't Granger-cause inflation if:

$$\phi_1^{\pi,u} = \phi_2^{\pi,u} = \phi_3^{\pi,u} = \phi_4^{\pi,u} = 0 \quad (4)$$

If unemployment does not Granger-cause inflation, then lagged values of unemployment are not useful for predicting inflation.

## 2.2 Granger-causality Robust Test

Suppose that the parameters in eq (2) are time varying, i.e. for  $j = 1, \dots, p$ , we replace  $\Phi_j$  with  $\Phi_{j,t}$ ,  $t = 1, \dots, T$ . Thus, eq (2) becomes:

$$y_{t+h} = \Phi_{1,t}y_{t-1} + \Phi_{2,t}y_{t-2} + \dots + \Phi_{p,t}y_{t-p} + \epsilon_{t+h} \quad (5)$$

Similarly, let  $\theta_t$  be a subset of  $vec(\Phi_{1,t}, \Phi_{2,t}, \dots, \Phi_{p,t})$ .

The null hypothesis of the Granger-causality robust test is:

$$H_0: \theta_t = 0 \quad \forall t = 1, 2, \dots, T \quad (6)$$

The statistics to test  $H_0$  in eq (6), following from Rossi (2005), are  $ExpW^*$  (the exponential Wald tests),  $MeanW^*$  (the mean Wald tests),  $Nyblom^*$  (the Nyblom test), and  $QLR^*$  (the Quandt likelihood ratio tests).<sup>2</sup>

The optimal exponential Wald test statistic ( $ExpW^*$ ) and the optimal mean Wald test statistic ( $MeanW^*$ ) are based on the exponential test statistics proposed in Andrews and Ploberger (1994). The optimal mean Wald test statistic is designed for alternatives that are very close to the null hypothesis; while the optimal exponential Wald test statistic is designed for testing against more distant alternatives. The optimal Nyblom test statistic ( $Nyblom^*$ ) is based on the Nyblom (1989) test, which is the locally most powerful invariant test for the constancy of the parameter process against the alternative that the parameters follow a random walk process. The optimal Quandt likelihood ratio test statistic ( $QLR^*$ ) is based on Andrews (1993) Sup-LR test (or the Quandt likelihood ratio (QLR) test), which considers the supremum of the statistics over all possible break dates of the Chow statistic designed for the alternatives for a fixed break date.

### 3 The gcr robustvar command

#### 3.1 The gcr robustvar command

##### Syntax

The `gcr robustvar` command is the Stata command that implements the VAR-based Granger-causality robust test. The general syntax of the `gcr robustvar` command is

```
gcr robustvar depvarlist, pos(#, #) [nocons horizon(#) lags(numlist) trimming(level)]
```

*depvarlist* is a list of dependent variables, that is, all the variables in  $y_t$  in the notation in eq (2).

*pos*(#, #) is a numeric list (i.e. "numlist" in Stata) including two integers indicating the positions of the targeted dependent variable and restricted regressor respectively. For example, if we are testing

---

<sup>2</sup>See Rossi (2005) for more details of constructing the statistics.

whether the second variable  $y_{2,t}$  Granger-causes the first variable  $y_{1,t}$  in the presence of instabilities, then we assign the numeric list as  $pos(1,2)$ , where the first interger 1 refers to the position of the targeted dependent variable in the VAR (i.e.  $y_{1,t}$  in this example) and the second interger 2 refers to the position of the targeted restricted regressor in the VAR (i.e.  $y_{2,t}$  in this example).

## Options

*noncons* suppresses the constant term. The default regression includes the constant term.

*horizon(#)* specifies the targeted horizon, i.e.  $h$  in the notation in eq (5). The default, i.e. not specifying *horizon(#)*, refers to a reduced-form VAR assuming homoskedastic idiosyncratic shocks. When *horizon(h)* ( $h \geq 0$ ) is specified, the command assumes heteroskedastic and serially correlated idiosyncratic shocks, and chooses the truncation lag used in the estimation of the long run variance. The truncation lag is automatically determined using Newey and West (1994) optimal lag-selection algorithm. Note that *horizon(0)* refers to a reduced-form VAR assuming heteroskedastic and serially correlated idiosyncratic shocks, and *horizon(h)* ( $h > 0$ ) refers to the  $(h + 1)$ -step-ahead forecasting model, see eq (5). For example, in a one-year-ahead VAR-LP forecasting model with quarterly data, *horizon(3)* should be specified.

*lags(numlist)* is a numeric list that specifies the lags included in the VAR. The default is *lags(1 2)*. This option takes a numlist and not simply an integer for the maximum lag. For example, *lags(2)* would include only the second lag in the model, whereas *lags(1 2)* would include both the first and second lags in the model. The shorthand to indicate the range follows "numlist" in Stata.

*trimming(level)* is the trimming parameter. As is standard in the structural break literature, the possible break dates are usually trimmed to exclude the beginning and end of the sample period. If we specify *trimming( $\mu$ )*, the range where we search for instabilities is set to be  $[\mu T, (1 - \mu)T]$ , where  $T$  is the number of total periods. The default is *trimming(0.15)*, which is recommended in the structural break literature and commonly used in practice.

## Stored results

`gcrobustvar` stores the following macros and matrices in `r()`:

Macros

r(cmd)	gcr robustvar
r(cmdline)	command as typed

### Matrices

r(stat)	A 4-by-1 matrix containing four statistics: $ExpW^*$ , $MeanW^*$ , $Nyblom^*$ , $SupLR^*$ .
r(pv)	A 4-by-1 matrix containing four p-values, corresponding respectively to $ExpW^*$ , $MeanW^*$ , $Nyblom^*$ , $SupLR^*$ .

## 3.2 Empirical Example of Practical Implementation in Stata

In what follows, we illustrate how to use the `gcr robustvar` command to implement the Granger-causality robust test in Stata. The data (`GCdata.xlsx`, provided with the article files) include quarterly U.S. data on the rate of price inflation ( $\pi_t$ ), the unemployment rate ( $u_t$ ), the interest rate ( $R_t$ , specifically, the federal funds rate) from 1959:I - 2000:IV. These are the same variables used in Stock and Watson (2001). Inflation is computed as  $\pi_t = 400 \times \ln(P_t/P_{t-1})$ , where  $P_t$  is the chain-weighted GDP price index. Quarterly data on  $u_t$  and  $R_t$  are quarterly averages of their monthly values.

Consider the inflation equation in (5):

$$\pi_t = c_t^\pi + \Phi_t^{\pi,\pi}(L)\pi_t + \Phi_t^{\pi,u}(L)u_t + \Phi_t^{\pi,R}(L)R_t + \epsilon_t^\pi \quad (7)$$

$$\text{where } \Phi_t^{\pi,\cdot}(L) = \phi_{1,t}^{\pi,\cdot}L + \phi_{2,t}^{\pi,\cdot}L^2 + \phi_{3,t}^{\pi,\cdot}L^3 + \phi_{4,t}^{\pi,\cdot}L^4$$

Suppose we are interested in testing whether  $R_t$  Granger-causes  $\pi_t$  and we want the test to be robust to instabilities over time. That is, we want to test whether the coefficients of lagged values of  $R_t$  are zero across time:

$$H_0 : \phi_{j,t}^{\pi,R} = 0 \quad \forall j = 1, 2, 3, 4 \quad \forall t = 1, 2, \dots, T$$

### **Implementing the Granger-causality Tests in the Presence of Instabilities**

The following scripts implement the Granger-causality robust test. We first import the data:

```

.
. * Import data
. import excel GCdata.xlsx, sheet(SW2001) cellrange(A1:C169) firstrow clear

. gen time = _n

. tsset time
    time variable:  time, 1 to 168
                  delta: 1 unit

```

Then we run the Granger-causality robust test using the `gcrobustvar` command. When we run the `gcrobustvar` command, important information (variables, lags, etc) will be displayed:

```

. gcrobustvar pi u R, pos(1,3) lags(1/4)
Running the Granger Causality Robust Test...
Setting:
Variables in VAR: pi u R
Lags in VAR:1 2 3 4
h is 0 (reduced-form VAR).
Trimming parameter is .15
Constant is included.
Assuming homoskedasticity in idiosyncratic shocks.

```

The results are displayed in the following script. The `gcrobustvar` command provides the four optimal test statistics ( $ExpW^*$ ,  $MeanW^*$ ,  $Nyblom^*$ ,  $QLR^*$ ) and their corresponding p-values.

```

Results of Granger Causality Robust Test: Lags of R Granger cause pi
ExpW*,MeanW*,Nyblom*,QLR* -- and their p-values below

              ExpW      MeanW      Nyblom      SupLR
statistics(pi:R) 11.838291 13.21676 5.7569834 32.111942
p-value(pi:R)   .01381882 .20035279 .03163363      0

```

Here is how we get all the inputs of the `gcrobustvar` command. `devarlist` lists the variables included in the VAR, i.e.  $\pi$ ,  $u$ ,  $R$  in this order. Since we are testing whether lags of the third variable  $R_t$  Granger-cause the first variable  $\pi_t$  in the presence of instabilities, we assign the following positions `pos(1,3)`. As for the options, we include the constant term and include four lags, i.e. `lags(1/4)`, as Stock and Watson (2001). Besides, we assume homoskedasticity and choose the standard trimming parameter 0.15.

Here is how to interpret the results. Let's take the exponential Wald tests statistics, denoted as  $ExpW^*$ , as an example. The value of the test statistic  $ExpW^*$  is 11.84, and the p-value is 0.01, which is smaller than the critical value at the 5% significance level. The test rejects the null hypothesis that

interest rate doesn't Granger-cause inflation for all  $t$  at the 5% significance level.

## 4 Comparison with the Traditional Granger-Causality Test

In this section, we compare the robust Granger-causality tests with the traditional Granger-causality test in the three-variable VAR model in Stock and Watson (2001). The VAR includes a constant term, four lags and assumes homoskedastic idiosyncratic shocks.

Table 1 reports the p-values of the traditional Granger-causality Wald statistics. The results show that  $\pi$  Granger-causes  $R$ ,  $u$  Granger-causes both  $\pi$  and  $R$ , and  $R$  Granger-causes  $u$  at the 5% significance level.

Table 1: Traditional Reduced-form VAR-based Granger-Causality Tests

Restricted Regressors	Dependent Variable		
	$\pi$	$u$	$R$
$\pi$	0.00	0.25	0.00
$u$	0.01	0.00	0.00
$R$	0.22	0.00	0.00

Note: This table reports p-values of the Wald statistics of the traditional Granger-causality test.  $h = 0$  (i.e. the reduced-form VAR model),  $lags = (1, 2, 3, 4)$ , assuming homoskedastic idiosyncratic shocks.

Table 2 reports the p-values of the robust Granger-causality test statistics (for  $ExpW^*$ ,  $MeanW^*$ ,  $Nyblom^*$  and  $QLR^*$ , respectively). We are testing whether the restricted regressor Granger-causes the dependent variable in the presence of instabilities. For example, if we consider the dependent variable  $\pi$  and the restricted regressor  $R$ , we are testing whether  $R$  Granger-causes  $\pi$  in a way robust to instabilities across time, i.e. whether the coefficients of lags of  $R$  are constant and equal to zero over time. The p-value of the  $ExpW^*$  statistics in Panel A in Table 2 is 0.01, so the test does reject the null at the 0.05% significance level. Hence,  $R$  does Granger-cause  $\pi$ . Comparing Table 1 and Table 2, the empirical conclusions differ if a researcher uses the Granger-causality robust test instead of the traditional Granger-causality test. In fact,  $R$  doesn't Granger-cause  $\pi$  at the 5% significance level in the traditional Granger-causality test, but  $R$  does Granger-cause  $\pi$  at the 5% significance level in the Granger-causality robust test according to the  $ExpW^*$ ,  $Nyblom^*$ ,  $SupLR^*$  test statistics. Hence, there is empirical evidence that lagged values of  $R$  can predict  $\pi$  but the predictive ability only shows up sporadically over time, which

is the reason why the traditional Granger-causality test doesn't detect it.

Table 2: Robust Granger-Causality Tests in the Reduced-form VAR

Panel A ExpW*			
	Dependent Variable		
Restricted Regressors	$\pi$	$u$	$R$
$\pi$	0.00	0.20	0.00
$u$	0.07	0.00	0.00
$R$	0.01	0.00	0.00

  

Panel B MeanW*			
	Dependent Variable		
Restricted Regressors	$\pi$	$u$	$R$
$\pi$	0.00	0.44	0.00
$u$	0.06	0.00	0.00
$R$	0.20	0.01	0.00

  

Panel C Nyblom*			
	Dependent Variable		
Restricted Regressors	$\pi$	$u$	$R$
$\pi$	0.00	0.22	0.00
$u$	0.08	0.00	0.00
$R$	0.03	0.02	0.00

  

Panel D QLR*			
	Dependent Variable		
Restricted Regressors	$\pi$	$u$	$R$
$\pi$	0.00	0.08	0.00
$u$	0.07	0.00	0.00
$R$	0.00	0.00	0.00

Note: This table reports p-values of the statistics of the Granger-causality robust test.  $h = 0$  (i.e. the reduced-form VAR model),  $lags = (1, 2, 3, 4)$ ,  $pistart = 0.15$ , assuming homoskedastic idiosyncratic shocks.

## 5 Robust Granger-Causality Tests in Local Projections

Section 4 considers the reduced-form VAR assuming homoskedastic idiosyncratic shocks. In this section, we extend the VAR analysis to Jorda's (2005) Local Projections by implementing the direct multistep

VAR-LP forecasting model in eq (2) and assuming heteroskedastic and serially correlated idiosyncratic shocks. Allowing for heteroskedasticity and serial correlation in idiosyncratic shocks is important when the researcher extends the VAR analysis to Local Projections, where the error terms in eq (2) can be both heteroskedastic and serially correlated.

We consider the one-year-ahead VAR-LP forecasting model with a constant term, four lags and assuming heteroskedastic and serially correlated idiosyncratic shocks. The setting is similar to Section 4 except that we specify  $h = 3$  and relax the homoskedasticity assumption.

The following is the command to implement the Granger-causality robust test to investigate whether the coefficients on  $R_{t-1}, R_{t-2}, R_{t-3}, R_{t-4}$  in the equation where the dependent variable is  $\pi_{t+3}$  are zero across time in the one-year-ahead VAR-LP forecasting model. To test other coefficients, the command is similarly implemented, except for adjusting the input of  $pos(\#, \#)$ .

```
. gcrobustvar pi u R, pos(1,3) lags(1/4) horizon(3)
Running the Granger Causality Robust Test...
Setting:
Variables in VAR: pi u R
Lags in VAR:1 2 3 4
h is 3 (4-step-ahead VAR-LP forecasting model).
Trimming parameter is .15
Constant is included.
Assuming heteroskedasticity and serial correlation in idiosyncratic shocks.
```

Table 3 reports the p-values of the robust Granger-causality test statistics (the  $ExpW^*$ ,  $MeanW^*$ ,  $Nyblom^*$  and  $QLR^*$  statistics respectively). The results show that lags of inflation ( $\pi$ ) can significantly forecast the one-year-ahead unemployment ( $u$ ) and interest rate ( $R$ ), lags of unemployment can significantly forecast the one-year-ahead inflation and interest rate, and lags of interest rate can significantly forecast the one-year-ahead inflation and unemployment.

Table 3: Robust Granger-causality Tests in the Direct Multistep VAR-LP Forecasting Model

Panel A ExpW*			
	Dependent Variable		
Restricted Regressors	$\pi$	$u$	$R$
$\pi$	0.00	0.00	0.00
$u$	0.00	0.00	0.00
$R$	0.00	0.00	0.00

  

Panel B MeanW*			
	Dependent Variable		
Restricted Regressors	$\pi$	$u$	$R$
$\pi$	0.00	0.00	0.00
$u$	0.00	0.00	0.00
$R$	0.00	0.00	0.00

  

Panel C Nyblom*			
	Dependent Variable		
Restricted Regressors	$\pi$	$u$	$R$
$\pi$	0.00	0.00	0.00
$u$	0.00	0.00	0.00
$R$	0.00	0.00	0.00

  

Panel D QLR*			
	Dependent Variable		
Restricted Regressors	$\pi$	$u$	$R$
$\pi$	0.00	0.00	0.00
$u$	0.00	0.00	0.00
$R$	0.00	0.00	0.00

Note: This table reports p-values of the statistics of the Granger-causality robust test.  $h = 3$  (i.e. the one-year-ahead VAR-LP forecasting model),  $lags = (1, 2, 3, 4)$ ,  $pistart = 0.15$ , assuming heteroskedastic and serially correlated idiosyncratic shocks.

## 6 References

Andrews, D.W., 1993. Tests for parameter instability and structural change with unknown change point. *Econometrica: Journal of the Econometric Society*, pp.821-856.

- Andrews, D.W. and Ploberger, W., 1994. Optimal tests when a nuisance parameter is present only under the alternative. *Econometrica: Journal of the Econometric Society*, pp.1383-1414.
- Boivin, J. and Giannoni, M.P., 2006. Has monetary policy become more effective?. *The Review of Economics and Statistics*, 88(3), pp.445-462.
- Clark, T.E. and McCracken, M.W., 2006. The predictive content of the output gap for inflation: resolving in-sample and out-of-sample evidence. *Journal of Money, Credit and Banking*, pp.1127-1148.
- Cogley, T. and Sargent, T.J., 2001. Evolving post-world war II US inflation dynamics. *NBER macroeconomics annual*, 16, pp.331-373.
- Cogley, T. and Sargent, T.J., 2005. Drifts and volatilities: monetary policies and outcomes in the post WWII US. *Review of Economic dynamics*, 8(2), pp.262-302.
- Granger, C.W., 1969. Investigating causal relations by econometric models and cross-spectral methods. *Econometrica: Journal of the Econometric Society*, pp.424-438.
- Jordà, O., 2005. Estimation and inference of impulse responses by local projections. *American economic review*, 95(1), pp.161-182.
- Kozicki, S. and Tinsley, P.A., 2001. Shifting endpoints in the term structure of interest rates. *Journal of monetary Economics*, 47(3), pp.613-652.
- Nyblom, J., 1989. Testing for the constancy of parameters over time. *Journal of the American Statistical Association*, 84(405), pp.223-230.
- Quandt, R.E., 1960. Tests of the hypothesis that a linear regression system obeys two separate regimes. *Journal of the American statistical Association*, 55(290), pp.324-330.
- Rossi, B., 2005. Optimal tests for nested model selection with underlying parameter instability. *Econometric theory*, 21(5), pp.962-990.
- Rossi, B., 2013. *Advances in Forecasting under Model Instability*. In: G. Elliott and A. Timmermann (eds.), *Handbook of Economic Forecasting, Volume 2B*, Elsevier Publications), 1203-1324.
- Sims, C.A., 1980. Macroeconomics and reality. *Econometrica: Journal of the Econometric Society*, pp.1-48.

- Stock, J.H. and Watson, M.W., 1996. Evidence on structural instability in macroeconomic time series relations. *Journal of Business & Economic Statistics*, 14(1), pp.11-30.
- Stock, J.H. and Watson, M.W., 1999. Forecasting inflation. *Journal of Monetary Economics*, 44(2), pp.293-335.
- Stock, J.H. and Watson, M.W., 2001. Vector autoregressions. *Journal of Economic perspectives*, 15(4), pp.101-115.
- Stock, J.H. and W Watson, M., 2003. Forecasting output and inflation: The role of asset prices. *Journal of Economic Literature*, 41(3), pp.788-829.
- Stock, J.H. and Watson, M.W., 2006. Forecasting with many predictors. *Handbook of economic forecasting*, 1, pp.515-554.